



Determinants of Breast Cancer screening: Application of Protection Motivation Theory

Fazlollah Ghofranipour ¹, Fatemeh Pourhaji ^{1,2,3,*}, Mohammad Hossein Delshad ^{1,2,3} and Fahime Pourhaji ⁴

¹Department of Health Education and Health Promotion, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

²Department of Public Health, School of Health, Torbat Heydariyeh University of Medical Sciences, Torbat Heydariyeh, Iran

³Health Sciences Research Center, Torbat Heydariyeh University of Medical Sciences, Torbat Heydariyeh, Iran

⁴Department of Health Education and Health Promotion, School of Health, Mashhad University of Medical Sciences, Mashhad, Iran

*Corresponding author: Department of Health Education and Health Promotion, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran. Email: fatemeh.pourhaji@modares.ac.ir

Received 2020 February 02; Revised 2020 April 20; Accepted 2020 May 02.

Abstract

Background: Breast cancer is a major public health problem in both developed and developing countries. The mortality rates of this disease are due to the lack of awareness about screening methods and late detection of breast cancer, which is high in Iran.

Objectives: The aim of this study was to determine breast self-examination (BSE) behaviors applying protection motivation theory (PMT).

Methods: In this cross-sectional study, the multi-stage cluster sampling method was used to 410 women aged 40 - 65 years old in Tehran, Iran. The questionnaire was completed through self-reported for each of the participants. PMT theoretical variables and BSE behavior are the basis of the data collection procedure. All analyses were performed using SPSS 20 for the windows. One-way ANOVA, chi-Square test, Independent Samples *t*-test, logistic regression, and Pearson correlation coefficient were applied. We set 0.05 as a criterion for statistical significance.

Results: The results indicated there were significant and positive correlations between the knowledge about breast cancer and self-efficacy of practicing BSE ($r = 0.43, P < 0.001$), response efficacy ($r = 0.20, P < 0.001$), and protection motivation ($r = 0.25, P = 0.003$).

Conclusions: Healthcare providers may consider PMT as a framework for developing educational interventions aiming at improving women's BSE behavior.

Keywords: Breast Self-Examination, Breast Cancer, Protection Motivation Theory

1. Background

Breast cancer disease is the most common cancer in the developed and developing countries according to the literature. Breast cancer is a major health problem in Iran (1, 2) and according to the latest national databases, age-standardized rate for breast cancer is 33.21 per 100,000 (2).

A recent study of breast cancer reported that the mean age for breast cancer in Iranian women is 5 years earlier compared to women from the developed countries (3) and increasing the trend of incidence is reported by the national cancer registry of project. Also, breast cancer is the fifth leading cause of cancer deaths and it is estimated 14.2% of death (2). Previous research performed in Iran showed that breast cancer has a significant impact on women's life (4).

Lack of awareness regarding risk factors, breast cancer screening methods, cultural taboos, feeling ashamed

to talk about breast cancer, lead to late detection and development of breast cancer and death. The easiest early detection that women could do is breast self-examination (BSE) (5).

BSE is a simple, effective, and helpful method of breast cancer screening, which is appropriate for all women. In addition, it is inexpensive, and increases self-awareness (6). There is no evidence on the effect of screening through BSE and no supportive role for BSE in the early detection of breast cancer. While according to the Kotka Pilot Project, BSE has better detection and decreases mortality, however, Swedish, Russian, and Shanghai studies demonstrated no improvements in mortality decreasing (7).

However, it has been shown that BSE may be of special importance in countries where breast cancer is a rising problem, but where mammography services are almost absent (8). There is a lack of a population-based mammo-

gram screening program in Iran. Therefore, it appears that BSE may be considered an attainable approach to empower women in the early detection of breast cancer.

BSE is helpful to women who have no access to other screening such as mammography. Despite the benefits of BSE, many women are inactive and different research has reported that the performance of BSE is low (9). The research suggested screening behavior is related to risk perception, benefit, barriers and reasoning processes that include personal and social factors and attitudes. Further studies are needed to understand the factors associated with the onset and maintenance of BSE throughout life.

The protection motivation theory (PMT) is a useful and social cognitive model for motivating individuals to use protective behavior. The PMT is frequently applied to breast cancer screening (10). According to the PMT model, women who have a higher perceived risk, and are susceptible to breast cancer and those who consider themselves at risk and serious disease would be more likely to affect regular BSE. Several studies suggested the effectiveness of PMT in Breast cancer screening (11-15). Although breast cancer is one of the few detected cancers in the initial phase, this level is very low in Iran.

2. Objectives

This study aimed to determine breast self-examination (BSE) behaviors applying protection motivation theory (PMT).

3. Methods

This cross-sectional study was done over a period of 6 months in health centers from May to October 2017. The statistical population of the study included 410 women aged 40 - 65 years old in Tehran, Iran. According to the rule of 5 - 10 individuals for each item (63×6), the sample size of 378 computer users was estimated. However, for more accuracy, the sample size was increased to 410 individuals (16). The multi-stage cluster sampling was implemented. First, the North, East, and Shemiranat networks were selected from 10 health networks of Shahid Beheshti University of Medical Sciences, Iran (SBUMS), and from each health network ($n = 3$), five urban health centers were randomly selected. Then from each health center ($n = 15$), 27 women over 40 years were randomly selected. Consequently, the sample size was computed at $27 \times 15 \approx 410$.

Women who met the following were eligible due to the inclusion/exclusion criteria. The exclusion criteria were a history or current diagnosis of cancer, pregnancy, and breastfeeding. Additionally, the following exclusion criteria were considered: insufficient knowledge of the Persian

language and insufficient physical and mental health to fill in the questionnaire. However, if someone suffered from any defect or disease interfering, they were excluded from the study. The demographic characteristics are presented by group in Table 1.

Table 1. Demographic Characteristics of Women

Variables	Mean \pm SD or No. (%)
Age	
40 - 50	43.69 \pm 2.7
51 - 60	53.35 \pm 2.2
Upper 60	61.9 \pm 2.4
Marital status	
Married	352 (88)
Single	13 (3.3)
Widow	26 (6.5)
Divorced	9 (2.2)
Women occupation	
Housewives	368 (92)
Working	32 (8)
Education level	
Primary	149 (37.3)
Secondary	117 (29.3)
More than high school	134 (32.9)

The research instrument included a self-reported questionnaire with 63 items. The questionnaire contained items of demographic information (4 items); knowledge questionnaire (20 items), PMT of breast cancer (37 items) and questions regarding BSE practice (2 items). The answers were 'true', 'false', and 'don't know'. The knowledge score was computed correct answers for all 20 questions (17). The PMT scale was developed in 1984 by McRae (18). In this study, first, a questionnaire was developed based on PMT and existing literature (12, 15, 19-27).

This questionnaire after a careful review and cultural adaptation and a few changes were made. The reliability and validity of the instrument were confirmed in the Iranian population (28). The final version of the questioner was average content validity ($s\text{-CVI}/\text{Ave}$) 0.80 that indicating adequate content validity. The reliability was determined by calculating the Cronbach's alpha coefficient. The coefficient α for the instrument was as follows: knowledge = 0.81, perceived vulnerability = 0.83, perceived severity = 0.79, fear = 0.84, response efficacy = 0.76, response cost = 0.84, self-efficacy = 0.84, perceived rewards = 0.83, protection motivation = 0.90.

BSE behavior was assessed as self-reported in BSE

within the past 1 month and the intended screening with the next month. The two following items assessed past behavior: "Have you done BSE in the last month?" (Yes/No.), and "How often have you done BSE in the last six months?" along with a seven-point scale (never once a month). Items with higher average scores + more frequency of past BSE behavior were collected (27).

All analyses were performed using SPSS 20 for the windows. One-way ANOVA, chi-square test, independent samples *t*-test, logistic regression, and Pearson correlation coefficient were applied. We set 0.05 as a criterion for statistical significance.

3.1. Ethics

The Ethics Committee of Tarbiat Modares University approved the study (IR.TMU.REC.1395.328) all participants gave written informed consent. Institutional ethics approval (IRCT2017061134472N1) was obtained before the start of the research.

4. Results

Four hundred participants responded in this study with a response rate of 97.5%. Ten questionnaires were excluded (four questionnaires because of lack of participation, three questionnaires because of uncompleted filling, three questionnaires because of having breast problems). The mean age of women was 45.69 ± 5.5 years. The results showed 91.25% of women did not have enough knowledge about breast cancer (score < 12). Most of the women were married (88%) (Table 1).

The results of the Pearson correlation coefficient test showed that there was no significant relationship between age and perceived severity ($r = 0.06$, $P = 0.17$), perceived vulnerability ($r = 0.07$, $P = 0.12$), response efficacy ($r = 0.01$, $P = 0.7$), response cost ($r = 0.09$, $P = 0.06$), perceived rewards ($r = 0.01$, $P = 0.7$), and self-efficacy ($r = 0.02$, $P = 0.63$), respectively. While there was a significant relationship between age and fear ($r = 0.141$, $P = 0.005$). The results of the One-way ANOVA test showed that there was a significant difference between the marital status and the mean score of perceived severity and vulnerability, response efficacy and self-efficacy so that married people had higher scores ($P \leq 0.05$). There was no significant relationship between the marital status and response cost, and perceived rewards ($P \geq 0.05$). Also, the results of the One-way ANOVA test showed that there was a significant difference between the education level and the mean score of perceived severity and vulnerability and response efficacy, and self-efficacy ($P \leq 0.05$), so women with higher education had a higher score in these structures.

According to logistic regression analysis; participants having knowledge on BSE practice were 3.46 times more likely to practice BSE [OR = 3.46, 95% CI (0.53 - 1.98)] compared with those less knowledgeable. Participants having a positive history of breast cancer in their family/friends were 2.59 times more likely to practice BSE [OR = 2.59, 95% CI (0.26 - 1.59)] compared with those having a negative history of breast cancer in their family/friends (Table 2).

Table 2. Logistic Regression Analysis Between the History of Breast Cancer in Family/Friends and Knowledge on BSE with BSE Practice^a

Variables	BSE Practice, No. (%)		OR (95% CI)
	Yes (N = 80)	No (N = 320)	
History of breast cancer in family/friends			
Positive	6 (7.5)	10 (3.12)	2.59 (0.26 - 1.59) ^b
Negative	74 (92.5)	310 (96.88)	1
Knowledge of BSE			
Knowledgeable	15 (18.8)	20 (6.2)	3.46 (0.53 - 1.98) ^b
Less knowledgeable	65 (81.2)	300 (93.8)	1

Abbreviation: BSE, breast self-examination

^aP value ≤ 0.05

^bReference category

The results of simple bivariate correlation showed there was significant and positive correlations between the knowledge about breast cancer and self-efficacy in practicing BSE ($r = 0.43$, $P < 0.001$), response efficacy ($r = 0.20$, $P < 0.001$), and protection motivation ($r = 0.25$, $P = 0.003$) (Table 3).

Table 3. Correlation Between Knowledge About Breast Cancer and Sub-Scale of PMT

Scale	Knowledge on Breast Cancer	
	r	P Value
Perceived vulnerability	0.10	0.1
Perceived severity	0.19	0.3
Fear	0.06	0.7
Response efficacy	0.20	< 0.001
Response cost	-0.21	< 0.001
Perceived rewards	-0.32	0.03
Self-efficacy	0.43	< 0.001
Protection motivation	0.25	0.003

Abbreviation: PMT, protection motivation theory

The finding suggested there were significant negative correlations between the response cost and knowledge of

breast cancer ($r = -0.21$, $P < 0.001$). Also, the results indicated there were significant negative correlations between perceived rewards and knowledge of breast cancer ($r = -0.32$, $P = 0.03$).

Furthermore, the results suggested there were no significant correlations between perceived vulnerability and severity with BSE practice. The results indicated there were positive and significantly correlated between the fear, response efficacy, self-efficacy, and protection motivation with BSE behavior so that means these structures were more in the practice group than the non-practice group ($P < 0.001$). While there were negative and significantly correlated between the response cost, the perceived rewards with BSE behavior means that these structures were less in the practice group than the non-practice group ($P < 0.001$) (Table 4).

Table 4. The Mean of the PMT Subscale on Breast Self-Examination

Scale	Mean \pm SD		P Value ^a
	BSE Practice (80)	Non BSE Practice (320)	
Perceived vulnerability	9.75 \pm 2.99	9.57 \pm 3.05	0.5
Perceived severity	5.50 \pm 2.22	5.82 \pm 2.12	0.2
Fear	15.70 \pm 7.2	12.95 \pm 6.97	0.001
Response efficacy	15.92 \pm 3.10	17.46 \pm 2.62	< 0.001
Response cost	21.33 \pm 7.2	23.02 \pm 6.38	0.041
Perceived rewards	8.5 \pm 2.83	9.29 \pm 2.54	0.01
Self-efficacy	12.10 \pm 1.1	9.2 \pm 2.8	< 0.001
Protection motivation	2.64 \pm 0.48	2.19 \pm 0.80	< 0.001

Abbreviation: BSE, breast self-examination, PMT, protection motivation theory
^aIndependent samples *t*-test

5. Discussion

The primary aim of the current study was to determine breast self-examination (BSE) behaviors applying protection motivation theory (PMT). In our sample, breast cancer awareness was low that it leads to late in referring to health centers. Due to the findings, only 14.5% of the participant ever heard about BSE, 8.7% had enough knowledge about breast cancer, and 91.25% had poor knowledge, while Baena-Canada et al. (29) reported that enough knowledge was 9.7% and in the study of Yadegarfar et al. (30) low knowledge was reported 55.7%. World Health Organization (WHO) recommends promoting knowledge and encouraging in the community through early detection and diagno-

sis of breast cancer in all women, especially women aged 40-69 years old who are returned health care centers or hospitals (31).

In many findings, BSE behavior was determined by the knowledge of women or having information on diagnostic methods of breast cancer (32). In our study, it found the knowledge is a significant variable in BSE. Similarly, in Hyun's research, it was suggested that women who are experienced to carry out BSE have a better level of knowledge of breast cancer (33). Breast cancer occurs in Iranian women earlier (34), so to enhance awareness about breast cancer screening can help in deducting mortality. It seems lifestyle changes and socioeconomic have a positive relationship with breast cancer (35). In this study, participants who had a positive history of breast cancer in their family/friends were more likely to practice BSE compared with those had a negative history. Similarly, some studies suggested this finding (36, 37). A positive history can act as a trigger that drives a person to take up a given breast cancer prevention behavior.

In this research, only 20% of the females reported doing regularly and monthly BSE. Similarly, some studies have indicated less than half of the participants really practice BSE monthly (38, 39). In the study of Mekuria et al. (32) performance of BSE was reported 13.4% and in the study of Badakhsh et al. (40) it was reported between 2.6% to 84.7% and an average 21.9%. While in the Didarloo et al. study, 24.6% (41) and in the study of Ertem and Kocer (42), 52% of women practiced BSE. In our study BSE performance was reported lower than the average. It seems one of the reasons this finding was lower literacy of women.

In the present study, the main source of information was the health care team. This indicates that health workers are effective. So that in the systematic review of Bouya et al. (43) it was reported that the most important sources of information were the healthcare team and it was confirmed in similar research. Nearly 21% of women have obtained information on breast cancer from TV/radio. Also, education based on the internet and social networks bring awareness of women effectively. In the current study, one of the sources of information was the Internet (26.5%). Tortolero-Luna et al. (44) described cancer information-seeking behavior and they reported that the Internet was the frequent sources of information about cancer (28.1%). The results showed there was no significant relationship between BSE practice and demographic variables (except for educational level). Similar to our study, in Jirojwong et al.'s study (45) and Dundar et al.'s research (46), it was explained that socio-demographic variables were not effective in BSE practice.

Similarly, the results of studies carried out by Fry and Prentice-Dunn (15), Boer and Seydel (47), Floyd et al. (19),

Hodgkins and Orbell (23), suggested the PMT is a beneficial framework to recognize factors that influence BSE for Iranian women. In this study perceived vulnerability and severity were not significant in explaining the BSE practice on a regular basis, but increased self-efficacy, response efficacy, and reduced BSE response cost and perceived rewards were significantly associated with BSE behavior. In Jordanian and U.S (48), Turkish (46), and Chinese (49) studies the perceived severity of women was reported as a non-significant predictor of BSE. It seems perceived severity is not a good predictor for breast cancer because this disease may be perceived by all women as an important and serious event, affecting the psychological, physical, and social aspects of life (46). Protection motivation was found to be a significant factor for BSE practice parallel to the results of Vahedian Shahroodi et al. (12) and Lee Champion (17).

Response efficacy was a significant variable predicting BSE practice. According to the finding of American studies, women who have more perceived response efficacy in BSE were more likely to practice BSE behavior (50, 51).

In our study self-efficacy was a significant factor for BSE practice and women who performed BSE on a regular basis had higher self-efficacy levels than non-practitioners women. The other studies indicated the various degree relationships between self-efficacy and breast cancer screening (52-54). In these studies, women who reported more self-efficacy in BSE were more likely to practice BSE regularly. In the current study, women who reported more fear of breast cancer were more likely to practice BSE regularly. Chen and Yang (55) reported similar results. Various studies have described the behavior of fear with both inhibitory and stimulating effects (56).

The present study has several strengths. This was one of the theory-driven studies examining breast cancer. Our findings provide evidence for the use of PMT in breast cancer prevention, which can be used as a framework for educational interventions in the field of breast cancer. Our study had several limitations. First, it was a cross-sectional design, so causal conclusions cannot draw. The sample of the research was middle-aged women in an urban area in Tehran, which does not necessarily reflect what happens among women in rural areas. So, the results of the study cannot be generalized to a larger population in Iran. Furthermore, the data were collected by a self-reported questionnaire, which may be a source of bias. Further studies with adequate confirmation of self-reported information built into their design are recommended.

5.1. Conclusion

Overall, the findings of our study suggest health care providers may consider PMT as a framework for developing

educational interventions aimed at improving women's BSE behavior.

Footnotes

Authors' Contribution: FP searched the data base and design, collected data and drafted the manuscript. FP, FGH contributed to conception and design of the study and revised the manuscript for intellectual content. MHD and FP designed the study, interpreted data, and revised the manuscript for important intellectual content. All authors read and approved the final manuscript.

Clinical Trial Registration Code: Institutional ethics approval (IRCT2017061134472N1) was obtained before the start of the research.

Conflict of Interests: There are no conflicts of interest.

Ethical Approval: The Ethics Committee of Tarbiat Modares University approved the study (IR.TMU.REC.1395.328) all participants gave informed written consent.

Funding/Support: This study was supported by Tarbiat Modares University.

References

- Ahmadi A, Ramazani R, Rezaghali T, Yavari P. Incidence pattern and spatial analysis of breast cancer in Iranian women: Geographical Information System applications. *Eastern Mediterranean Health Journal*. 2018;**24**(4).
- Nafissi N, Khayamzadeh M, Zeinali Z, Pazooki D, Hosseini M, Akbari ME. Epidemiology and Histopathology of Breast Cancer in Iran versus Other Middle Eastern Countries. *Middle East Journal of Cancer*. 2018;**9**(3):243-51.
- Akbari A, Razzaghi Z, Homaei F, Khayamzadeh M, Movahedi M, Akbari ME. Parity and breastfeeding are preventive measures against breast cancer in Iranian women. *Breast Cancer*. 2011;**18**(1):51-5. doi: [10.1007/s12282-010-0203-z](https://doi.org/10.1007/s12282-010-0203-z).
- Pourhaji F, Ghofranipour F. Designing and Psychometric Evaluation of Breast Self-Examination Behavior Predicting Scale (BSEBPS). *Int J Cancer Manag*. 2018;**11**(12). e74266. doi: [10.5812/ijcm.74266](https://doi.org/10.5812/ijcm.74266).
- Zare N, Haem E, Lankarani KB, Heydari ST, Barooti E. Breast Cancer Risk Factors in a Defined Population: Weighted Logistic Regression Approach for Rare Events. *J Breast Cancer*. 2013;**16**(2):214-9.
- Coleman C. Early Detection and Screening for Breast Cancer. *Seminars in Oncology Nursing*. 2017;**33**(2):141-55. doi: [10.1016/j.soncn.2017.02.009](https://doi.org/10.1016/j.soncn.2017.02.009).
- Haji-Mahmoodi M, Montazeri A, Jarvandi S, Ebrahimi M, Haghighat S, Harirchi I. Breast Self-Examination: Knowledge, Attitudes, and Practices Among Female Health Care Workers in Tehran, Iran. *The breast journal*. 2002;**8**(4):222-5.
- Miller AB, Wall C, Baines CJ, To T. Canadian National Breast Screening Study-2: 13-Year Results of a Randomized Trial in Women Aged 50-59 Years. *JNCI: Journal of the National Cancer Institute*. 2000;**92**(18):1490-9. doi: [10.1093/jnci/92.18.1490](https://doi.org/10.1093/jnci/92.18.1490).
- Asghari E, Nahamin M, Khoshtarash M, Ghanbari A, Parizad N, Mahdavi N, et al. The relationship between health belief and breast self-examination among Iranian university students. *Int. J. Women Health Reprod. Sci*. 2016;**4**:110-3.

10. Meyerowitz BE, Chaiken S. The effect of message framing on breast self-examination attitudes, intentions, and behavior. *Journal of personality and social psychology*. 1987;**52**(3):500.
11. Rahaei Z, Ghofranipour F, Morowatisharifabad MA, Mohammadi E. Determinants of cancer early detection behaviors: application of protection motivation theory. *Health promotion perspectives*. 2015;**5**(2):138.
12. Vahedian Shahroodi M, Pourhaje F, Esmaily H. Investigating the effectiveness of protection motivation, perceived self-Efficacy and perceived response costs by behavior of breast self-examination. *The Iranian Journal of Obstetrics, Gynecology and Infertility*. 2013;**15**(40):1-9.
13. Helmes AW. Application of the Protection Motivation Theory to Genetic Testing for Breast Cancer Risk. *Preventive Medicine*. 2002;**35**(5):453-62. doi: [10.1006/pmed.2002.1110](https://doi.org/10.1006/pmed.2002.1110).
14. Karmakar M, Pinto SL, Jordan TR, Mohamed I, Holiday-Goodman M. Predicting Adherence to Aromatase Inhibitor Therapy among Breast Cancer Survivors: An Application of the Protection Motivation Theory. *Breast Cancer: Basic and Clinical Research*. 2017;**11**:1.1782234176945E+15. doi: [10.1177/1178223417694520](https://doi.org/10.1177/1178223417694520).
15. Fry RB, Prentice-Dunn S. Effects of a psychosocial intervention on breast self-examination attitudes and behaviors. *Health Education Research*. 2006;**21**(2):287-95. doi: [10.1093/her/cyh066](https://doi.org/10.1093/her/cyh066).
16. Hajizadeh E, Asghari M. Statistical methods and analyses in health and biosciences a research methodological approach. *Tehran: Jahade Daneshgahi Publications*. 2011;**395**.
17. Lee Champion V. Use of the health belief model in determining frequency of breast self-examination. *Research in Nursing & Health*. 1985;**8**(4):373-9. doi: [10.1002/nur.4770080410](https://doi.org/10.1002/nur.4770080410).
18. McCrae RR. Situational determinants of coping responses: Loss, threat, and challenge. *Journal of Personality and Social Psychology*. 1984;**46**(4):919-28. doi: [10.1037/0022-3514.46.4.919](https://doi.org/10.1037/0022-3514.46.4.919).
19. Floyd DL, Flournoy JM, Prentice-Dunn S. Effects of persuasive message order on coping with breast cancer information. *Health Education Research*. 2001;**16**(1):81-4. doi: [10.1093/her/16.1.81](https://doi.org/10.1093/her/16.1.81).
20. Rippetoe PA, Rogers RW. Effects of components of protection-motivation theory on adaptive and maladaptive coping with a health threat. *Journal of Personality and Social Psychology*. 1987;**52**(3):596-604. doi: [10.1037/0022-3514.52.3.596](https://doi.org/10.1037/0022-3514.52.3.596).
21. Lee JB. *Breast cancer fear, mammography fear, and mammography adherence of African American women in Bridgeport, Connecticut [dissertation]*. New York University; 2011.
22. Champion V, Maraj M, Hui S, Perkins AJ, Tierney W, Menon U, et al. Comparison of tailored interventions to increase mammography screening in nonadherent older women. *Preventive medicine*. 2003;**36**(2):150-8.
23. Hodgkins S, Orbell S. Can protection motivation theory predict behaviour? A longitudinal test exploring the role of previous behaviour. *Psychology & Health*. 1998;**13**(2):237-50. doi: [10.1080/08870449808406749](https://doi.org/10.1080/08870449808406749).
24. Smith SM, Ford JS, Rakowski W, Moskowitz CS, Diller L, Hudson MM, et al. Inconsistent mammography perceptions and practices among women at risk of breast cancer following a pediatric malignancy: a report from the Childhood Cancer Survivor Study. *Cancer Causes & Control*. 2010;**21**(10):1585-95.
25. Schwarzer R, Renner B. Social-cognitive predictors of health behavior: action self-efficacy and coping self-efficacy. *Health psychology*. 2000;**19**(5):487.
26. Ajzen I. The theory of planned behavior. *Organizational behavior and human decision processes*. 1991;**50**(2):179-211.
27. Prestwich A, Conner M, Lawton R, Bailey W, Litman J, Molyneux V. Individual and collaborative implementation intentions and the promotion of breast self-examination. *Psychology and Health*. 2005;**20**(6):743-60.
28. Pourhaji F, Ghofranipour F. *Evaluation of Educational Program based on Integrated model health process approach (HAPA) with Protection Motivation Theory (PMT) on breast cancer screening Behavior in upper 40 years of Tehran [dissertation]*. Tehran: School of Medical Sciences; 2019.
29. Baena-Cañada JM, Rosado-Varela P, Expósito-Álvarez I, González-Guerrero M, Nieto-Vera J, Benítez-Rodríguez E. Women's perceptions of breast cancer screening. Spanish screening programme survey. *The Breast*. 2014;**23**(6):883-8. doi: [10.1016/j.breast.2014.09.010](https://doi.org/10.1016/j.breast.2014.09.010).
30. Yadegarfar G, Salami F, Mostajeran M, Ansari R, Rejali M, Aghdak P. Factors Engaged With Women Performance in Doing Regular Breast Self-examination or Referring to Health Centers for Examination and Mammography. *Quarterly of Horizon of Medical Sciences*. 2018;**24**(1):53-60.
31. WHO. *Knowledge into Action Cancer Control WHO Guide for Effective Programmes*. WHO.; 2018. 5 p. Contract No.: 92 4 154733 8.
32. Mekuria M, Nigusse A, Tadele A. Breast Self-Examination Practice and Associated Factors Among Secondary School Female Teachers in Gammo Gofa Zone, Southern, Ethiopia. *Breast Cancer: Targets and Therapy*. 2020;**12**:1.
33. Lee E. Breast Self-Examination Performance Among Korean Nurses. *Journal for Nurses in Professional Development*. 2003;**19**(2):81-7.
34. Farhood B, Geraily G, Alizadeh A. Incidence and Mortality of Various Cancers in Iran and Compare to Other Countries: A Review Article. *Iranian journal of public health*. 2018;**47**(3):309-16. [PubMed: [29845017](https://pubmed.ncbi.nlm.nih.gov/29845017/)].
35. Esmaeizadeh N, Salahi-Moghaddam A, Khoshdel A. Geographic distribution of important cancers in Iran. *Bimonthly Journal of Hormozgan University of Medical Sciences*. 2015;**19**(2):66-76.
36. Juárez-García DM, Téllez A. The Health Belief Model and Prediction of Breast Self-examination Practices in Female Mexican College Students. *Psychology in Russia. State of the Art*. 2019;**12**(3).
37. Bertoni N, de Souza MC, Crocamo S, Szklo M, de Almeida LM. Is a Family History of the Breast Cancer Related to Women's Cancer Prevention Behaviors? *International Journal of Behavioral Medicine*. 2019;**26**(1):85-90. doi: [10.1007/s12529-018-9737-9](https://doi.org/10.1007/s12529-018-9737-9).
38. Fung S. Factors associated with breast self-examination behaviour among Chinese women in Hong Kong. *Patient Education and Counseling*. 1998;**33**(3):233-43. doi: [10.1016/S0738-3991\(98\)00023-8](https://doi.org/10.1016/S0738-3991(98)00023-8).
39. Alwan NA, Al-Diwan JK, Al-Attar WM, Eliessa RA. Knowledge, attitude & practice towards breast cancer & breast self examination in Kirkuk University, Iraq. *Asian Pacific Journal of Reproduction*. 2012;**1**(4):308-11. doi: [10.1016/S2305-0500\(13\)60098-6](https://doi.org/10.1016/S2305-0500(13)60098-6).
40. Badakhsh M, Balouchi A, Taheri S, Bouya S, Ahmadidarehsima S, Aminifard M. Attitude and Practice Regarding Breast Cancer Early Detection among Iranian Women: A Systematic Review. *Asian Pacific journal of cancer prevention : APJCP*. 2018;**19**(1):9-16. doi: [10.22034/APJCP.2018.19.1.9](https://doi.org/10.22034/APJCP.2018.19.1.9). [PubMed: [29373873](https://pubmed.ncbi.nlm.nih.gov/29373873/)].
41. Didarloo A, Nabilou B, Khalkhali HR. Psychosocial predictors of breast self-examination behavior among female students: an application of the health belief model using logistic regression. *BMC Public Health*. 2017;**17**(1):861. doi: [10.1186/s12889-017-4880-9](https://doi.org/10.1186/s12889-017-4880-9).
42. Ertem G, Kocer A. Breast self-examination among nurses and midwives in Odemis health district in Turkey. *Indian J Cancer*. 2009;**46**(3):208-13. doi: [10.4103/0019-509x.52955](https://doi.org/10.4103/0019-509x.52955). [PubMed: [19574672](https://pubmed.ncbi.nlm.nih.gov/19574672/)].
43. Bouya S, Balouchi A, Ahmadidarehsima S, Badakhsh M. Knowledge and Source of Information About Early Detection Techniques of Breast Cancer Among Women in Iran: A Systematic Review. *Journal of cancer prevention*. 2018;**23**(1):51-60. doi: [10.15430/JCP.2018.23.1.51](https://doi.org/10.15430/JCP.2018.23.1.51). [PubMed: [29629349](https://pubmed.ncbi.nlm.nih.gov/29629349/)].
44. Tortolero-Luna G, Rutten L, Hesse B, Davis T, Kornfeld J, Sanchez M, et al. Health and Cancer Information Seeking Practices and Preferences in Puerto Rico: Creating an Evidence Base for Cancer Communication Efforts. *J Health Commun*. 2010;**15** Suppl 3:30-45. doi: [10.1080/10810730.2010.522698](https://doi.org/10.1080/10810730.2010.522698).
45. Jirojwong S, MacLennan R. Health beliefs, perceived self-efficacy, and breast self-examination among Thai migrants in Brisbane. *Journal of Advanced Nursing*. 2003;**41**(3):241-9. doi: [10.1046/j.1365-2648.2003.02552.x](https://doi.org/10.1046/j.1365-2648.2003.02552.x).
46. Dündar PE, Özmen D, Öztürk B, Haspolat G, Akyıldız F, Çoban S, et al. The knowledge and attitudes of breast self-examination and mam-

- mography in a group of women in a rural area in western Turkey. *BMC Cancer*. 2006;**6**(1):43. doi: [10.1186/1471-2407-6-43](https://doi.org/10.1186/1471-2407-6-43).
47. Boer H, Seydel ER. *Protection Motivation Theory*. In Conner M, Norman P, editors, *Predicting Health Behaviour: Research and Practice with Social Cognition Models*. Eds. Mark Conner, Paul Norman. Buckingham: Open University Press; 1996. p. 95-120.
 48. Petro-Nustus W, Mikhail BI. Factors Associated with Breast Self-Examination Among Jordanian Women. *Public Health Nursing*. 2002;**19**(4):263-71. doi: [10.1046/j.1525-1446.2002.19406.x](https://doi.org/10.1046/j.1525-1446.2002.19406.x).
 49. Lu ZJ. Variables associated with breast self-examination among Chinese women. *Cancer nursing*. 1995;**18**(1):29-34. [PubMed: [7866974](https://pubmed.ncbi.nlm.nih.gov/7866974/)].
 50. Rutledge DN, Davis GT. Breast self-examination compliance and the health belief model. *Oncology nursing forum*. 1988;**15**(2):175-9. [PubMed: [3357830](https://pubmed.ncbi.nlm.nih.gov/3357830/)].
 51. Ayala GX. *Differences in the practice of breast self-examination by Latina & caucasian-American women: the role of acculturation, group orientation, locus of control and health beliefs*. 2015.
 52. Milne S, Sheeran P, Orbell S. Prediction and Intervention in Health-Related Behavior: A Meta-Analytic Review of Protection Motivation Theory. *Journal of Applied Social Psychology*. 2000;**30**(1):106-43. doi: [10.1111/j.1559-1816.2000.tb02308.x](https://doi.org/10.1111/j.1559-1816.2000.tb02308.x).
 53. Luszczynska A, Schwarzer R. Planning and Self-Efficacy in the Adoption and Maintenance of Breast Self-Examination: A Longitudinal Study on Self-Regulatory Cognitions. *Psychology & Health*. 2003;**18**(1):93-108. doi: [10.1080/0887044021000019358](https://doi.org/10.1080/0887044021000019358).
 54. KELLI GARCIA, TRACI MANN. From 'I Wish' to 'I Will': Social-cognitive Predictors of Behavioral Intentions. *Journal of Health Psychology*. 2003;**8**(3):1-15. doi: [10.1177/13591053030083005](https://doi.org/10.1177/13591053030083005).
 55. Chen L, Yang X. Using EPPM to Evaluate the Effectiveness of Fear Appeal Messages Across Different Media Outlets to Increase the Intention of Breast Self-Examination Among Chinese Women. *Health Communication*. 2018:1-8. doi: [10.1080/10410236.2018.1493416](https://doi.org/10.1080/10410236.2018.1493416).
 56. Gore TD, Bracken CC. Testing the Theoretical Design of a Health Risk Message: Reexamining the Major Tenets of the Extended Parallel Process Model. *Health Education & Behavior*. 2005;**32**(1):27-41. doi: [10.1177/1090198104266901](https://doi.org/10.1177/1090198104266901).